



When should my child return to school?

Chicken Pox When all blisters have dried over		Conjunctivitis None*		Diarrhoea & Vomiting 48 hours from last episode and well enough to return	Flu Until recovered	Hand, foot & mouth None*	German Measles 5 days from onset of rash	Glandular Fever None*
Hand, Foot & Mouth None*	Head Lice None*	Impetigo When lesions are crusted & healed or 48 hours after commencing antibiotics	Measles 4 days from onset of rash and well enough	Mumps 5 days from onset of swelling	Ringworm Not usually required but treatment is needed		Rubella (German measles) 5 days from the appearance of the rash	Scabies 24 hours after the first dose of treatment
Scarlet Fever 24 hours after commencing antibiotic treatment	Slapped Cheek None* Once rash has developed	Threadworms None* Treatment recommended	Tonsillitis None*	Whooping Cough 48 hours after commencing antibiotics	Respiratory Infections, including Corona virus (Covid-19) Children with mild symptoms such as a runny nose, sore throat, or mild cough, who are otherwise well - NONE Children who are unwell and have a high temperature should stay at home until they are well enough. If a child has a positive Covid-19 test result they should stay at home for 3 days after the day, they took the test.			

*No need to stay off school but school should be informed. [Link to Health & Security Agency https://www.gov.uk/government/publications/health-protection-in-schools-and-other-childcare-facilities/managing-specific-infectious-diseases-a-to-z](https://www.gov.uk/government/publications/health-protection-in-schools-and-other-childcare-facilities/managing-specific-infectious-diseases-a-to-z)